


Amendments to the Claims

Please cancel claims 10-27 without prejudice. Please add new claims 28-47 as shown below in the Listing of Claims.

Listing of Claims

1-27. (Cancelled)

28. (New) A process for the hydrogenation of a compound, comprising hydrogenating a C₆-C₁₈ aromatic substituted amino acid or C₆-C₁₈ aromatic substituted amino alcohol in the presence of a platinum-rhodium mixed catalyst, wherein said C₆-C₁₈ aromatic substituted amino acid or C₆-C₁₈ aromatic substituted amino alcohol has the general formula (I):



wherein

n is 0, 1 or 2;

R¹ is a (C₆-C₁₈) aryl, or a (C₇-C₁₉) aralkyl, wherein aryl groups are optionally substituted with halogen, (C₁-C₈) alkoxy, (C₁-C₈) acyl, or (C₁-C₈) acyloxy;

R² is H, OH, (C₁-C₈) alkyl, (C₂-C₈) alkoxyalkyl;

R³ and R⁴ are each independently H, a (C₁-C₈) alkyl, a (C₆-C₁₈) aryl, or together denote an =O function;

P¹ and P² are each independently hydrogen, an amino protective group or together stand for a bifunctional amino protective group;

P³ is hydrogen, a hydroxyl protective group, or a carboxyl protective group; and

the carbon atom marked with * is an asymmetrical carbon atom.

29. (New) The process of claim 28, wherein R² is H.

30. (New) The process of claim 29, wherein R^3 and R^4 are H, or together denote an =O function.
31. (New) The process of claim 28, wherein the ratio of platinum to rhodium in said platinum-rhodium mixed catalyst is between 20:1 and 1:1 (w/w).
32. (New) The process of claim 28, wherein said platinum-rhodium mixed catalyst is used in a quantity of 0.1 to 20 wt%, relative to the compound undergoing hydrogenation.
33. (New) The process of claim 28, wherein said platinum-rhodium mixed catalyst is adsorbed on a support.
34. (New) The process of claim 28, wherein said hydrogenation is performed in the presence of a solvent selected from the group consisting of: water; an alcohol; an ether; and mixtures thereof.
35. (New) The process of claim 28, wherein said hydrogenation is performed at a temperature of 10°C to 150°C.
36. (New) The process of claim 28, wherein said process comprises reacting said C_6 - C_{18} aromatic substituted amino acid or C_6 - C_{18} aromatic substituted amino alcohol with hydrogen gas in the presence of said platinum-rhodium mixed catalyst and under a hydrogen pressure of between 1 and 100 bar.
37. (New) The process of claim 28, wherein:
- a) R^2 is H;
 - b) R^3 and R^4 are H, or together denote an =O function; and
 - c) the ratio of platinum to rhodium in said platinum-rhodium mixed catalyst is between 20:1 and 1:1 (w/w).
38. (New) The process of claim 37, wherein said platinum-rhodium mixed catalyst is used in a quantity of 0.1 to 20 wt%, relative to the compound undergoing hydrogenation.

39. (New) The process of claim 38, wherein:
- a) said hydrogenation is performed in the presence of a solvent selected from the group consisting of: water; and an alcohol;
 - b) said hydrogenation is performed under a hydrogen pressure of between 1 and 100 bar; and
 - c) said hydrogenation is performed at a temperature of 10°C to 150°C.
40. (New) The process of claim 39, wherein said platinum-rhodium mixed catalyst is adsorbed on a support.
41. (New) A process for the hydrogenation of a compound selected from the group consisting of: L-phenylalanine, D-phenylalanine, L-phenylglycine, D-phenylglycine, L-tyrosine or D-tyrosine, comprising hydrogenating said compound in the presence of a platinum-rhodium mixed catalyst.
42. (New) The process of claim 41, wherein the ratio of platinum to rhodium in said platinum-rhodium mixed catalyst is between 20:1 and 1:1 (w/w).
43. (New) The process of claim 42, wherein said platinum-rhodium mixed catalyst is used in a quantity of 0.1 to 20 wt%, relative to the compound undergoing hydrogenation.
44. (New) The process of claim 43, wherein said hydrogenation is performed in the presence of a solvent selected from the group consisting of: water; an alcohol; an ether; and mixtures thereof.
45. (New) The process of claim 44, wherein said hydrogenation is performed at a temperature of 10°C to 150°C.
46. (New) The process of claim 45, wherein said process comprises reacting said C₆-C₁₈ aromatic substituted amino acid or C₆-C₁₈ aromatic substituted amino alcohol with hydrogen gas in the presence of said platinum-rhodium mixed catalyst and under a hydrogen pressure of between 1 and 100 bar.

47. (New) The process of claim 46, wherein said platinum-rhodium mixed catalyst is adsorbed on a support.